INDEX TO VOLUME XX

New names and the final members of new combinations are in **bold face type.**

Abies, 44, 45; amabilis, 301; balsamea, 44; concolor, 301; Fraseri, 241; grandis, 301; lasiocarpa, 301
Absidia caerulea, 176
Acacia acuifera, 215; amentacea, 215; coriophylla, 215; Farnesiana, 215; unijuga, 215; Wrightii, 215
Acanthonitschkea, 29
Acer, 315; glabrum, 102, 109, 112, 113; negundo, 242, 244; rubrum, 319; saccharinum, 241
Achlya americana, 172; DeBaryana, 172; gracilipes, 172; intermedia, 171; polyandra, 172; racemosa, 173
Adenoropium gossypifolium, 76, 80
Aecidium, 62, 75, 117; dominicanum, 80, 81; Phacae-frigidae, 42; Tournefortiae, 75, 82; Trillii, 117–119, 124
Agropyron spicatum, 110
Albugo Bliti, 173
Alisma brevipes, 166; Plantago, 167; subcordata, 167
Alnus, 107, 317; insignis, 102; tenuifolia, 102, 104, 107, 172
Amaranthus retroflexus, 173
Amorpha fruticosa, 235
Amyris balsamifera, 216; elemifera, 216; Plumieri, 215
Andropogon, 54; Sorghum sudanensis, 74, 80
Ansonia, 99
Anthostoma, 306, 307, 322, 323, 326, 329–333, 335, 336; gastrinum, 329; grandineum, 330
Anthostomella, 306, 307, 330, 332, 335; minor, 330
Anthurium acaule, 218; aripoense, 53; dominicense, 218; Holtonianum, 218; Isertianum, 218; panduratum, 218; scandens, 218; venosum, 218
Apocynum sibiricum, 245
Arachis hypogaea, 68, 80
Arbutus, 298; Menziesii, 298, 300
Arctostaphylos, 297, 298
Arcyria cinerea, 28; cinerea var. digitata, 28; decipiens, 112; denudata, 28, 102, 350
Armillaria mellea, 1
Arrangement of the spores in the ascus of Neurospora sitophila, The sexuality and, Marguerite S. Wilcox, 3
Arthur's rust work, Doctor, Fred J. Seaver, 115
Arthur, J. C., Notes on arctic Uredinales, 41
Artocarpus communis, 75, 80
Aschersonia, 53; Andropogonis, 54; disciformis, 53; parasitica, 54; viridans, 53; viridula, 53
Ascobolus, 9; carbonarius, 3, 10; furfuraceus, 4; immersus, 4
Ascomycetes—IV. Some Hypocreales from Trinidad, Studies in tropical, 52; V. Species of Phyllachora, Studies in tropical, Fred J. Seaver, 214
Ascophora elegans, 177
Ascopolyporus, 52, 53; polychrous, 52
Aster, 99
Ateleia cubensis, 216
Auerswaldia Miconiae, 222
Azotobacter, 266, 267, 271
Baccharis pilularis, 295
Bacillus amylovorus, 252; anthracis, 271; cereus, 265; Chauvoei, 271; coli, 255, 257, 270; mallei, 271; mesentericus, 257, 259, 264, 272; mycoides, 264; prodigiosus, 263; subtilis, 257, 259, 264, 265, 268, 272; tumefaciens, 257; typhi, 259, 272; typhosus, 259; vulgatus, 259, 272
Bacterial species—I. Morphologic variations, Variations within a, H. R. Rosen, 251
Bacterium fluorescens, 268; pneumoniae, 268
Badhamia capsulifera, 103, 348; nitens, 102, 112; populina, 103; utricularis, 103
Balansia, 142, 194, 211
Bambos vulgaris, 76, 80
Barrus, M. F., Trillium rust, 117
Basidiophora Kellermanii, 174
Bastardia viscosa, 70, 80
Begonia, 23
Betula, 106, 108, 111, 317; fontinalis, 102, 105, 106, 109, 169, 236; lenta, 238; nigra, 319; papyrifera, 110; populifolia, 236
Bihai Bihai, 70, 80

INDEX TO VOLUME XX

New names and the final members of new combinations are in **bold face type.**

Abies, 44, 45; amabilis, 301; balsamea, 44; concolor, 301; Fraseri, 241; grandis, 301; lasiocarpa, 301
Absidia caerulea, 176
Acacia acuifera, 215; amentacea, 215; coriophylla, 215; Farnesiana, 215; unijuga, 215; Wrightii, 215
Acanthonitschkea, 29
Acer, 315; glabrum, 102, 109, 112, 113; negundo, 242, 244; rubrum, 319; saccharinum, 241
Achlya americana, 172; DeBaryana, 172; gracilipes, 172; intermedia, 171; polyandra, 172; racemosa, 173
Adenoropium gossypifolium, 76, 80
Aecidium, 62, 75, 117; dominicanum, 80, 81; Phacae-frigidae, 42; Tournefortiae, 75, 82; Trillii, 117–119, 124
Agropyron spicatum, 110
Albugo Bliti, 173
Alisma brevipes, 166; Plantago, 167; subcordata, 167
Alnus, 107, 317; insignis, 102; tenuifolia, 102, 104, 107, 172
Amaranthus retroflexus, 173
Amorpha fruticosa, 235
Amyris balsamifera, 216; elemifera, 216; Plumieri, 215
Andropogon, 54; Sorghum sudanensis, 74, 80
Ansonia, 99
Anthostoma, 306, 307, 322, 323, 326, 329–333, 335, 336; gastrinum, 329; grandineum, 330
Anthostomella, 306, 307, 330, 332, 335; minor, 330
Anthurium acaule, 218; aripoense, 53; dominicense, 218; Holtonianum, 218; Isertianum, 218; panduratum, 218; scandens, 218; venosum, 218
Apocynum sibiricum, 245
Arachis hypogaea, 68, 80
Arbutus, 298; Menziesii, 298, 300
Arctostaphylos, 297, 298
Arcyria cinerea, 28; cinerea var. digitata, 28; decipiens, 112; denudata, 28, 102, 350
Armillaria mellea, 1
Arrangement of the spores in the ascus of Neurospora sitophila, The sexuality and, Marguerite S. Wilcox, 3
Arthur's rust work, Doctor, Fred J. Seaver, 115
Arthur, J. C., Notes on arctic Uredinales, 41
Artocarpus communis, 75, 80
Aschersonia, 53; Andropogonis, 54; disciformis, 53; parasitica, 54; viridans, 53; viridula, 53
Ascobolus, 9; carbonarius, 3, 10; furfuraceus, 4; immersus, 4
Ascomycetes—IV. Some Hypocreales from Trinidad, Studies in tropical, 52; V. Species of Phyllachora, Studies in tropical, Fred J. Seaver, 214
Ascophora elegans, 177
Ascopolyporus, 52, 53; polychrous, 52
Aster, 99
Ateleia cubensis, 216
Auerswaldia Miconiae, 222
Azotobacter, 266, 267, 271
Baccharis pilularis, 295
Bacillus amylovorus, 252; anthracis, 271; cereus, 265; Chauvoei, 271; coli, 255, 257, 270; mallei, 271; mesentericus, 257, 259, 264, 272; mycoides, 264; prodigiosus, 263; subtilis, 257, 259, 264, 265, 268, 272; tumefaciens, 257; typhi, 259, 272; typhosus, 259; vulgatus, 259, 272
Bacterial species—I. Morphologic variations, Variations within a, H. R. Rosen, 251
Bacterium fluorescens, 268; pneumoniae, 268
Badhamia capsulifera, 103, 348; nitens, 102, 112; populina, 103; utricularis, 103
Balansia, 142, 194, 211
Bambos vulgaris, 76, 80
Barrus, M. F., Trillium rust, 117
Basidiophora Kellermanii, 174
Bastardia viscosa, 70, 80
Begonia, 23
Betula, 106, 108, 111, 317; fontinalis, 102, 105, 106, 109, 169, 236; lenta, 238; nigra, 319; papyrifera, 110; populifolia, 236
Bihai Bihai, 70, 80

Biologic studies in the Sphaeriales—I, 187; II, Julian H. Miller, 305
Bisexual mycelia, Unisexual conidia from, B. O. Dodge, 226
Bixa orellana, 80
Blastocladia ramosa, 169
Boletus, 46, 250
Bolinia, 310, 334; tubulina, 310
Bonar, Lee, Studies on some California fungi, 292
Borreria laevis, 72, 80
Botryorhiza, 62; Hippocrateae, 65, 81
Botryosphaeria, 193, 198; Bakeriana, 193; Quercum, 193; Ribis, 193
Botrytis, 141, 145, 147, 152; cinerea, 141; effusa, 173; farinosa, 173; ganglioniformis, 175; Jonesii, 178; parasitica, 174
Bourreria succulenta, 216
Boyce, J. S., Lophodermium Abietis on Pseudotsuga taxifolia, 301
Brachyelytrum, 125; erectum, 119, 120, 122–124
Brachyramphus intybaceus, 79–81
Bradburya virginiana, 220
Brassica, 23; Rapa, 23
Bremia Lactucae, 175
British Guiana and Surinam, Myxomycetes from, Frank A. Gilbert, 27
Bruce Fink, Lichenologist, Robert B. Wylie, 1
Bryophyllum pinnatum, 23
Bullaria, 68–71
Byrsonima crassifolia, 63, 80

Cacao, 55, 56
Caeoma hibisciatum, 70
Caeomurus Halstedii, 125
Cajan Cajan, 66, 80
California fungi, Studies on some, Lee Bonar, 292
Calonectria flavida, 56
Calonyction, 98
Calostilbe, 248; longiasca, 248; **striispora,** 248
Camarops, 310, 333, 334, 336; hypoxyloides, 310
Camarosporium **Betulae,** 236
Camillea, 336
Campanula, 22, 98
Canna, 69, 80
Capparis Grisebachii, 217
Capriola dactylon, 69, 80

Cardiospermum microcarpum, 68, 80
Carpinus caroliniana, 326
Cassia crista, 66, 80; emarginata, 65; Fistula, 217; marilandica, 240
Castanea, 313, 315
Castanopsis chrysophylla, 294; sempervirens, 294
Catacauma Brittoniana, 216; Myrciae, 220
Ceanothus sanguineus, 237
Cenchrus echinatus, 69, 80; viridis, 69, 80
Ceratiomyxa fruticulosa, 27; fruticulosa var. flexuosa, 27
Cercomonas, 350; longicauda, 350
Cercospora canescens, 236
Cerotelium, 62; desmium, 64, 81; Fici, 64, 81
Chaetocladium Jonesii, 178
Chamaecyparis, 19
Chamaesyce hirta, 68, 80; hypericifolia, 68, 81; prostrata, 68, 81
Chaetosphaeria phaeostroma, 199, 200
Chenopodium, 23; album, 173, 241
Chloris ciliata, 68, 81; paraguayensis, 68, 81
Chlorosplenium, 130, 133, 135; lividum, 130–133
Chondrioderma quitense, 105
Chrysanthemum, 22
Chrysopsis, 98
Chytridium ampullaceum, 159; Euglenae, 165; globosum, 160; pollinis, 161; Saprolegniae, 159; vagans, 161
Ciboria, 127–133, 135–141, 143–146, 148, 155; amentacea, 128, 129, 131, 144; bolaris, 128, 144; Caucus, 128, 132, 144; ciliatospora, 128; firma, 128–130, 144; fructicola, 154; livida, 132; rhizophila, 128; tremellosa, 128
Cichorium Endivia, 23
Ciferri, R., A new genus of the subfamily Nitschkieae, 29
Cissus sicyoides, 64, 81
Cladochytrium, 167; Alfalfae, 167; Alismatis, 166, 167; **maculare,** 166
Cladophora Kuetzingiana, 168
Clasterisporium, 244
Clathrus denudatus, 102
Clavicens, 191, 192, 194, 211
Clostridium butyricum, 271
Coccolobis laurifolia, 224; uvifera, 75, 81
Cogswellia, 174; leptocarpa, 174; montana, 174
Coleosporium, 62, 76, 97, 98; apocynaceum, 99; Campanulae, 98; carneum, 98; delicatulum, 99; domingensis, 77, 82; Elephantopodis, 62, 81, 98; Helianthi, 99; inconspicuum, 99; Ipomoeae, 98; Lacinariae, 98; Madiae, 99; minutum, 100; Plumierae, 77; ribicola, 99; Senecionis, 99; Solidaginis, 99; Sonchi-arvensis, 98; Terebinthinaceae, 99

Coleosporium occurring in the United States and a list of the host species, A key to the known aecial forms of, George G. Hedgcock, 97
Coleus Blumei, 23
Collema pulposum, 4
Colletotrichum carpophilum, 242; **Viciae,** 242
Collybia dryophila, 31, 33
Collybia dryophila in pure culture, The fruiting of, R. F. Poole, 31
Comandra Richardsiana, 36
Comatricha flaccida, 103; typhoides, 28, 103, 104
Commelina, 23
Contributions to our knowledge of western Montana fungi—I. Myxomycetes, 101; II. Phycomycetes, Paul W. Graff, 158
Convolvulus, 98
Cook, W. R. Ivimey, and Miss E. M. Holt, Some observations on the germination of the spores of some species of Mycetozoa, 340
Coprinus Lagopus, 3
Cordyceps, 192, 194
Coreopsis, 22, 29
Cornutia pyramidata, 75, 81
Corylus, 317, 321
Coryneum, 243; microstictum, 243; Negundinis, 244; septosporioides, 244
Craterium minutum, 104
Creosphaeria, 330
Cribaria purpurea, 104; rufa, 104
Cronartium, 38; Comandrae, 36–38; ribicola, 37
Cronartium Comandrae and Melampsora Medusae, The uredinia of, E. H. Moss, 36
Crossopsora, 62; notata, 63, 80
Croton angustatus, 63, 81; flavens, 224; lucidus, 224
Cryptosporium, 245; **acicolum,** 244; **Boycei,** 245
Cucurbita Pepo, 23
Curcas Curcas, 76, 81
Cylindrosporium Ceanothi, 237; Shepherdiae, 238; **sibiricum,** 245
Cyperus ferax, 68, 81
Cystopus Bliti, 173
Cystothyrium, 240
Cytospora, 295

Dahlia, 22
Daldinia, 306, 310, 328, 329, 332, 333, 335, 336; concentrica, 307, 328, 329; Eschscholzii, 328, 329; vernicosa, 328, 329
Daphne, 22
Dark-field microscopy in the study of fungi, Edwin E. Honey and W. R. Fisher, 88
Darluca filum, 125
Dearness, John, New and noteworthy fungi—V, 235
Delphinium, 22
Desmella, 62; superficialis, 66, 81
Diachaea leucopoda, 348
Diatrype, 307, 308, 324; disciformis, 189; stigma, 314
Dibotryon morbosum, 199, 201
Dicaeoma, 68–76, 79; Majanthae, 125
Dichromena ciliata, 69, 81; colorata, 69
Diderma brunneolum, 108; radiatum, 104
Didymium anomalum, 105; difforme, 105; nigripes, 28; quitense, 105; tigrinum, 107
Dinemasporium **corrugatum,** 240
Diplanes saprolegnioides, 171
Disaeta, 299; **Arbuti,** 299
Doassansia Alismatis, 167
Doctor Arthur's rust work, Fred J. Seaver, 115
Dodge, B. O., Spore formation in asci with fewer than eight spores, 18; Unisexual conidia from bisexual mycelia, 226
Dolichos multiflorus, 23
Dothichloe, 194
Dothidea, 208; collecta, 199–201; inclusa, 219; Lathyri, 220; Myrciae, 220; Tragiae, 224
Dothidella **castanicola,** 294; Castanopsidis, 294
Dothidina peribebuyensis, 222
Dothiora, 198
Draparnaldia plumosa, 166
Dryopteris subtetragona, 66, 81
Drypetes glauca, 218; lateriflora, 218
Duranta repens, 219

Echinochloa frumentacea, 23
Eichornia crassipes, 81
Elephantopus, 98; mollis, 62, 81
Eleutheranthera ruderalis, 73, 81
Elsota virgata, 224
Empusa Muscae, 178
Endophyllum, 62; circumscriptum, 64, 81; Stachytarphetae, 64, 82
Endoxylon, 322
Enteridium Rozeanum, 348
Entophlyctis, 162
Entyloma Meliloti, 247
Erodium cicutarium, 175
Erwin F. Smith, Frederick V. Rand, 181
Eucalyptus, 239, 297, 298; globulus, 297
Euglena, 166; viridis, 165, 166.

Eupatorium, 77, 81
Euphorbia, 39
Euthamia, 100

Fagus, 313
Fern rust new to the United States, An interesting, H. W. Thurston, Jr., 44
Ficus Carica, 64, 81; subscabrida, 217
Filix fragilis, 45
Fink, Lichenologist, Bruce, Robert B, Wylie, 1
Fisher, W. R., and Edwin E. Honey, Dark-field microscopy in the study of fungi, 88
Fitzpatrickia, 29; **Massae,** 30
Fomes pinicola, 285; roseus, 276, 281–290
Fomes roseus and Trametes subrosea, Temperature and moisture relations of, Walter H. Snell, W. G. Hutchinson and K. H. N. Newton, 276
Fraxinus, 315; oregana, 296
Fruiting of Collybia dryophila in pure culture, The, R. F. Poole, 31
Fuligo cinerea, 105; intermedia, 105; septica, 345, 346, 348, 351
Fulminaria, 163; Hedenii, 163; mucophila, 165
Funastrum clausum, 74, 81
Fungi of Santo Domingo—II. Uredinales, Frank D. Kern, 60

Galactia, 220
Galactinia succosa, 4
Gaya occidentalis, 70, 81
Geoglossum glabrum, 4
Geopyxis Rapulum, 132
Geranium viscosissimum, 175
Germination of the spores of some species of Mycetozoa, Some observations on the, W. R. Ivimey Cook and Miss E. M. Holt, 340
Gilbert, Frank A., Myxomycetes from British Guiana and Surinam, 27
Gloeosporium Chenopodii, 241; **melleum,** 241; **multipunctatum,** 241
Gossypium, 64; barbadense, 64, 81
Gouania lupuloides, 70, 71, 81; polygama, 70, 81
Graff, Paul W., Contributions to our knowledge of western Montana fungi—I. Myxomycetes, 101; II. Phycomycetes, 158
Grossularia, 99
Guignardia, 195, 196, 198, 208; Bidwellii, 199, 200
Gymnosporangium, 19

Hamelia erecta, 76, 81
Harknessia Arctostaphyli, 297, 298; Eucalypti, 297; uromycoides, 297, 298
Harpochytrium, 163; Hedenii, 163, 165; Hyalothecae, 163, 165; intermedium, 165
Hedgcock, George G., A key to the known aecial forms of Coleosporium occurring in the United States and a list of the host species, 97
Helianthus, 99
Hemitrichia clavata, 112, 350; Vesparium, 106, 348
Hendersonula **pinicola,** 236
Heracleum lanatum, 239
Hercospora, 199
Heterotrichum cymosum, 223
Hippocratea volubilis, 65, 81
Holcus halepensis, 74, 81; Sorghum, 74, 80, 81
Holt, Miss E. M., and W. R. Ivimey Cook, Some observations on the germination of the spores of some species of Mycetozoa, 340
Honey, Edwin E., The monilioid species of Sclerotinia, 127; and W. R. Fisher, Dark-field microscopy in the study of fungi, 88
Hordeum sativum, 23
Humaria granulata, 4; rutilans, 4
Hutchinson, W. G., K. H. N. Newton and Walter H. Snell, Temperature and moisture relations of Fomes roseus and Trametes subrosea, 276
Hyaloceros, 299
Hyalopsora, 38; Polypodii, 45
Hydnobolites sp., 4
Hydnum, 250
Hymenoscypha, 132; Candolleana, 132; ciborioides, 133; Curreyana, 133; Duriaeana, 133; filipes, 133; sclerotiorum, 132; tuberosa, 132
Hypocrea, 53, 57, 307, 308; amazonica, 53; viridans, 54
Hypocreales from Trinidad, Studies in tropical Ascomycetes—IV. Some, Fred J. Seaver, 52
Hypocrella, 52–54, 194, 211; Andropogonis, 54; viridans, 53
Hypoderma, 301; Hedgcockii, 240; lineare, 236
Hypodermella, 301; Laricis, 240
Hyponectria Phaseoli, 223
Hypoxylon, 83, 195, 196, 199, 202, 306–312, 314, 315, 318, 319, 321, 323–337; annulatum, 84, 85, 312, 318–320, 327, 332; atropunctatum, 324; atropurpureum, 84, 85; atrorufum, 84; bifrons, 84, 86; Blakei,

84; callostroma, 84, 86; caries, 84, 87; Catalpae, 84, 86; coccineum, 83–85, 192, 203, 308, 311–313, 317, 329, 332; cohaerens, 84, 85, 311; cclliculosum, 85; concentricum, 308; confluens, 312, 321, 327, 332; decorticatum, 85, 86; durissimum, 85, 86; effusum, 312, 319, 320; enteromelum, 84, 85; exsurgens, 332; fragiforme, 84; fuscopurpureum, 315; fuscum, 308, 312, 315, 317; glomeratum, 308; granulosum, 85, 308; gregale, 312, 320; Howeianum, 84, 85, 199, 202, 206, 311, 312, 313, 315–317, 328, 332, illitum, 312, 320; leucocreas, 332; luteum, 310; Malleolus, 312, 320; marginatum, 84–86, 312, 315, 318–320; Michelianum, 312, 320; Morsei, 84, 312, 320, 321; multiforme, 83–85, 308, 317; nummularium, 308, 310; perforatum, 84–86, 315, 316; Petersii, 332; placentiforme, 328, 329, 332; polyspermum, 319; rubiginosum, 83, 86, 311, 312, 315–317, 326, 334; Sassafras, 84, 86; semiimmersum, 312, 321, 331, 332; serpens, 84, 85, 87, 312, 320, 321; sphinctericum, 307; stigmateum, 324; suborbiculare, 326; transversum, 85; turbinulatum, 84; udum, 311, 321, 322, 330, 332; ustulatum, 308, 309, 328
Hypoxylon, Notes on the synonymy of some species of, C. L. Shear, 83
Hyptis capitata, 71, 81; suaveolens, 73, 81

Impatiens, 175; aurella, 175; Sultani, 23
Indigofera mucronata, 65, 81; suffruticosa, 65, 81
Inga Inga, 65, 81
Interesting fern rust new to the United States, An, H. W. Thurston, Jr., 44
Ipomoea, 80, 81, 98
Isandrina emarginata, 65, 81
Isoachlya monilifera, 172
Iva xanthiifolia, 174

Jacquemontia nodiflora, 66, 81
Jacquinia, 219, 220; Berterii, 219
Jambos Jambos, 74, 81
Jasminum grandiflorum, 78; pubescens, 78, 81
Jatropha gossypifolia, 76

Kabatia, 292
Keithia, 19, 20; Chamaecyparissi, 19; thujina, 20
Kern, Frank D., Fungi of Santo Domingo—II. Uredinales, 60
Key to the known aecial forms of Coleosporium occurring in the United States and a list of the host species, A, George G. Hedgcock, 97
Kretzschmaria, 336
Kuehneola, 62; Fici, 64; Gossypii, 64; malvicola, 66, 82
Kusanoopsis, 208

Laboulbenia, 18; chaetophora, 4
Labrella, 292
Lachnea scutellata, 4; stercorea, 4
Lacinaria, 98
Lactuca intybacea, 79, 81; Ludoviciana, 175; virosa, 175
Lagenidium, 168, 256; globosum, 168; Rabenhorstii, 169
Lamproderma columbinum, 106; robustum, 106; Sauteri, 106, 107; violaceum, 107
Lamprospora, 47; areolata, 47; Maireana, 47; tuberculata, 47; tuberculatella, 47
Lantana involucrata, 72, 81
Larix occidentalis, 103, 104, 108, 112, 240
Lars Romell, C. L. Shear, 49
Lasiacis divaricata, 67, 81
Lasiobotrys, 292; affinis, 292, 293, 294; Lonicerae, 292
Leersia oryzoides, 124; oryzoides var. japonica, 124; virginica, 124
Lentinus lepideus, 280; tigrinus, 280
Lenzites sepiaria, 277, 278, 279
Leonotis, 72; nepetaefolia, 72, 81
Lepargyrea canadensis, 238
Lepidoderma tigrinum, 107
Leptomassaria, 330
Leptosphaeria, 198, 205, 206, 209; acuta, 210; Doliolum, 201, 206, 208–210
Leptostroma **Hedgcockii,** 240
Leptostromella **Cassiae,** 240
Leptothyrella, 240; **Laricis,** 240
Leptothyrium, 292; **Pseudotsugae,** 239; **stenosporum,** 239
Leucostoma **Sequoiae,** 295
Libertella **ulmicola,** 246; Ulmi-suberosae, 246
Ligusticum, 42
Lindera, 87
Liquidambar, 86
Liriodendron tulipifera, 242
Lithocarpus densiflorus, 300
Lonchocarpus domingensis, 65, 81
Lonicera, 292, 293; hispidula, 292
Lophodermium, 301; Abietis, 301, 302

Lophodermium Abietis on Pseudotsuga taxifolia, J. S. Boyce, 301
Lupinus micranthus, 297; succulentus, 297
Lycogala epidendrum, 28, 107, 340, 345, 346, 350, 351
Lycoperdon epidendrum, 107; radiatum, 104; vesparium, 106
Lychnis alba, 237

Macbridella, 55, 248; striispora, 54, 55, 248
Madia, 99
Malvastrum corchorifolium, 73, 81; coromandelianum, 73, 81
Marssonia **Sonchi,** 243
Massaria, 198
Massarina, 198
Mayepea domingensis, 220
? Mazzantia fennica, 220
Medicago, 167; sativa, 174
Meibomia cana, 67, 81; scorpiurus, 67, 81; tortuosa, 73, 81
Melampsora, 38, 39; arctica, 42, 43; Bigelowii, 42; Helioscopiae, 39; Laricis-capraearum, 39; Lini, 39; Medusae, 36, 38, 39; populina, 39
Melampsora Medusae, The uredinia of Cronartium Comandrae and, E. H. Moss, 36
Melanconis, 198, 199
Melanomma, 198; seminis, 295
Melanospora, 192, 211, 306, 335
Melanthera, 66, 81
Melilotus alba, 23; indica, 247
Mesosphaerum capitatum, 71, 81; suaveolens, 73, 82
Miconia laevigata, 223; prasina, 223; Sintenisii, 223
Micropolis, 58
Micropuccinia, 68, 70, 72–75, 79
Milesia pycnograndis, 44
Milesina, 44
Miller, Julian H., Biologic studies in the Sphaeriales—I, 187; II, 305
Mimosa Ceratonia, 65, 82
Monilia, 152, 153; sitophila, 3, 226
Monilinia, 127, 152, 153, 155; **fructicola,** 153, 155
Monilioid species of Sclerotinia, The, Edwin E. Honey, 127
Monochaeta, 299; **pinicola,** 244
Montana fungi—I. Myxomycetes, Contributions to our knowledge of western, 101; II. Phycomycetes, Contributions to our knowledge of western, Paul W. Graff, 158
Morchella, 8; bispora, 18; esculenta, 4
Morphologic variations, Variations within a bacterial species—I., H. R. Rosen, 251
Morus alba, 240
Moss, E. H., The uredinia of Cronartium Comandrae and Melampsora Medusae, 36
Mougeotia, 159; sphaerocarpa, 161
Mucor abundans 176, 178; circinelloides, 176, 177; elegans, 177; hiemalis, 176; Mucedo, 176, 178; plumbeus 176, 177; racemosus, 177; sphaerocephalus, 176; spinescens, 176, 177; spinosus, 176; stolonifer, 177
Mycetozoa, Some observations on the germination of the spores of some species of, W. R. Ivimey Cook and Miss E. M. Holt, 340
Mycodiplosis, 125
Mycosphaerella, 198; Fragariae, 200
Myiophyton Cohnii, 178
Myriangium, 204, 207–209; Duriaei, 208, 209
Myrica paniculata, 221
Myriogenospora, 194, 211
Myriophyllum, 23
Myxomycetes, Contributions to our knowledge of western Montana fungi—I., Paul W. Graff, 101
Myxomycetes from British Guiana and Surinam, Frank A. Gilbert, 27
Myxosporium, 242; **megallanto,** 241; **negundicolum,** 242; **roseum,** 242
Myzocytium, 168, 256; proliferum, 168

Naegeliella Reinschii, 170
Nectandra patens, 221
Nectria **indusiata,** 58; rhytidospora, 56; setosa, 57; striispora, 55; suffulta, 57
Neottiella albocincta, 4
Neurospora, 202, 226, 227, 335; crassa, 3, 15, 226; sitophila, 3, 5–12, 14, 15, 19, 226, 231; tetrasperma, 3, 4, 6–8, 19, 20, 226, 227, 230, 231, 233, 234
Neurospora sitophila, The sexuality and arrangement of the spores in the ascus of, Marguerite S. Wilcox, 3
New and noteworthy fungi—V, John Dearness, 235
New genus of the subfamily Nitschkieae, A, R. Ciferri, 29
Newton, K. H. N., Walter H. Snell and W. G. Hutchinson, Temperature and moisture relations of Fomes roseus and Trametes subrosea, 276
Nigredo, 66–68; digitata, 125; Halstedii, 125
Nitschkia, 29

Nitschkieae, A new genus of the subfamily, R. Ciferri, 29
Nodulosphaeria, 206
Notes on arctic Uredinales, J. C. Arthur, 41
Notes on the parisitology of Sclerotium Rolfsii, Ruth Davis Paintin, 22
Notes on the synonymy of some species of Hypoxylon, C. L. Shear, 83
Nummularia, 85, 192, 306–308, 310, 312, 323–326, 332, 334, 335; atropunctatum, 312; Bulliardii, 189, 310, 312, 319, 320, 322, 324; clypeus, 312, 315, 323, 325; discreta, 85, 310, 312, 322, 326, 330, 333; dryophila, 310; hypophloea, 325; lutea, 310; microplaca, 325; punctulata, 312, 325, 326; tinctor, 325

Observations on the germination of the spores of some species of Mycetozoa, Some, W. R. Ivimey Cook and Miss E. M. Holt, 340
Ocotea leucoxylon, 222
Oedogonium crenulato-costatum, 159; plagiostomum, 160; plusiosporum, 169
Oidium, 231, 232
Oligostroma acicola, 245
Olpidiopsis Saprolegniae, 159
Olpidium pendulum, 158
Ombrophila, 133, 135–137; strobilina, 133, 135
Ontotelium digitatum, 125
Ophiobolus graminis, 4
Ophiodothis, 194, 211
Ornithogalum umbellatum, 23
Osmorrhiza longistylis, 168
Otidea aurantia, 4
Oxalis, 23

Paintin, Ruth Davis, Notes on the parasitology of Sclerotium Rolfsii, 22
Panicum barbinode, 67, 82; fasciculatum, 73; fasciculum, 82; maximum, 67, 73, 82
Parasitology of Sclerotium Rolfsii, Notes on the, Ruth Davis Paintin, 22
Parsonsia Parsonsia, 76, 82
Parthenium, 99
Pasania, 300
Paspalum conjugatum, 75, 82; Lindenianum, 69, 82; plicatulum, 75, 82
Pavonia coccinea, 66, 82
Penicillium, 146
Penzigia, 334
Perichaena chrysosperma, 28
Peridermium, 44, 97; acicolum, 99; apocynaceum, 99; californicum, 99; carneum, 98; delicatulum, 99; Elephantopodis, 98; Fischeri, 98; floridanum, 98; fragile, 98; Helianthi, 99; inconspicuum, 99; Ipomoeae, 98; minutum, 100; montanum, 99; oblongisporium, 99; pycnograndis, 44; ribicola, 99; Rostrupii, 98; Terebinthinaceae, 99; Weirii, 100
Peronospora effusa, 173; gangliformis, 175; Geranii, 175; Kellermanii, 174; obducens, 175; parasitica, 174; ribicola, 175; Trifoliorum, 174; Viciae, 174
Persea gratissima, 219; Persea, 219
Pestalozzia, 299; Castagnei, 300
Petroselinum hortense, 23
Peziza, 128, 129, 146; minuta, 104; Sclerotiorum, 129; Stevensoniana, 4; tuberosa, 129; vesiculosa, 4, 6
Phaca, 43
Phakopsora, 62; Crotonis, 63; **dominicana,** 63, 81; fenestrala, 78, 82; Meibomiae, 78, 81; mexicana, 63
Phaseolus, 223; multiflorus, 23; vulgaris, 236
Philocopra coeruleotecta, 4
Philodendron, 218
Phleospora, 237
Phlyctidium, 161
Phlyctochytrium, 162; equale, 160, 162; stellatum, 162
Phoma, 295; samararum, 296, 297; thermopsidicola, 295
Phycomyces nitens, 226
Phycomycetes, Contributions to our knowledge of western Montana fungi—II., Paul W. Graff, 158
Phyllachora, 214, 217; Acaciae, 215; **amyridicola,** 215; **Amyridis,** 215; Ateleiae, 216; Bourreriae, 216; **Brittoniana,** 216; Canafistulae, 217; **conspicua,** 217; drypeticola, 218; Engleri, 218; fusicarpa, 219; gratissima, 219; inclusa, 219; Lathyri, 220; Mayepeae, 220; Myricae, 220; Nectandrae, 221; nitens, 221; ocoteicola, 222; **Pennellii,** 222; peribebuyensis, 222; Phaseoli, 223; Philodendronis, 218; Randiae, 223; Securidacae, 224; simplex, 224; texana, 215; Tragiae, 224; Zanthoxyli, 224; zanthoxylicola, 225
Phyllachora, Studies in tropical Ascomycetes—V. Species of, Fred J. Seaver, 214
Phyllactinia, 6; corylea, 4, 18

Phyllanthus grandifolius, 78, 82
Phyllosticta castanicola, 294; innumera, 296; **Lupini,** 297; **sparsa,** 296; viridis, 296
Physalospora Phaseoli, 223
Physarella oblonga, 28
Physarum auriscalpium, 107; bitectum, 108, 110; bogoriense, 27; brunneolum, 108; carneum, 108; columbinum, 106; Diderma, 108, 109; didermoides, 349; oblatum, 108; polycephalum, 27, 109; sinuosum, 108, 109; tenerum, 27, 110; testaceum, 109
Physoderma, 167; maculare, 166, 167
Physopella, 75
Phytophthora, 281
Piaropus crassipes, 80–82
Picea Engelmanni, 111; excelsa, 301; sitchensis, 301
Pinus apacheca, 100; Banksiana, 100; canariensis, 100; caribaea, 100; chihuahuana, 100; clausa, 100; contorta, 100; Coulteri, 100; echinata, 100, 244; edulis, 100; Elliotti; 100; glabra, 100, 244; heterophylla, 100; insignis, 100; Laricio, 100, Laricio austriaca, 100; Mayriana, 100; Murrayana, 100; nigra austriaca, 100; nigra Poiretiana, 100; palustris, 100, 244; pinea, 100; ponderosa, 100, 106, 107, 158; ponderosa scopulorum, 100; pungens, 100; radiata, 100; resinosa, 100; rigida, 100, 240; Sabiniana, 100; serotina, 100; Strobus, 236, 240; sylvestris, 100; taeda, 100, 244; Thunbergii, 100; virginiana, 100, 237, 244
Piptocephalis Freseniana, 177
Plasmopara Geranii, 175; obducens, 175; ribicola, 175
Plectodiscella, 204
Pleomassaria, 198
Pleospora, 198, 205, 209; herbarum, 206, 209
Pleurocystis Fresenii, 177
Pluchea purpurascens, 79, 82
Plumiera rubra, 77, 82
Podocrea, 57
Podocrella, 57; **poronioides,** 57
Podospora anserina, 18
Polygonatum, 139; biflorum, 138; multiflorum, 138; pubescens, 138, 139; vulgare, 138
Polyphagus, 166; Euglenae, 165
Polypodium vulgare, 44
Polyporus, 52; abietinus, 279; benzoinus, 279, 280; pargamenus, 279; resinosus, 279, 280
Poole, R. F., The fruiting of Collybia dryophila in pure culture, 31
Populus, 39, 151; alba, 128; tremula, 128; tremuloides, 109, 243; trichocarpa, 103, 105, 111, 112, 169, 172
Poronia, 57, 332, 334, 336; punctata, 204
Porostigme, 196
Prinus, 87
Priva lappulacea, 72, 82
Prospodium, 62, 63; Amphilophii, 63; appendiculatum, 63, 66, 82; bahamense, 63; plagiopus, 63; **Tabebuiae,** 63, 82
Proteus vulgaris, 259, 272
Protomyces macularis, 166, 167
Prunus spinosa, 78, 82
Pseudolpidium Saprolegniae, 159
Pseudoperisporium, 196
Pseudosphaeria, 198, 204, 205
Pseudotsuga mucronata, 107, 110–112, 161, 170; taxifolia, 239, 245, 301, 302
Pseudotsuga taxifolia, Lophodermium Abietis on, J. S. Boyce, 301
Pseudovalsa, 199
Pteris longifolia, 23
Puccinia, 62, 70; Arachidis, 68, 80; Arechavelatae, 68, 80; Arenariae, 41; Arthurella, 79; 80; cacabata, 68, 81; canaliculata, 68, 81; Cannae, 69, 80; Cardamines-bellidifoliae, 42; Cenchri, 69, 80; Chaetochloae, 69, 82; Chloridis, 68; Cochleariae, 42; Cruciferarum, 42; Cynodontis, 69, 80; Dichromenae, 69, 81; dominicana, 72; Drabae, 42; fuscella, 69, 82; Gouaniae, 70, 81; Heliconiae, 70, 80; heterospora, 70, 80–82; hibisciata, 70, 82; Holboellii, 41; Hyptidis, 71, 81; impedita, 71, 82; inflata, 71, 82; insueta, 71; insulana, 71, 82; invaginata, 71, 81; Lantanae, 72, 81, 82; lateritia, 72, 80; Leonotidis, 72, 81; levis, 73, 79, 82; Malvacearum, 73, 81; medellinensis, 73, 81; Melampodii, 73, 81, 82; obliqua, 74, 81; pallescens, 74, 82; Panici, 73; Plucheae, 79, 82; Polygoni-vivipari, 42; polysora, 79, 82; proximella, 79; Pruni-spinosae, 78; Psidii, 74, 81; purpurea, 74, 81; Rivinae, 74, 82; scleriicola, 75, 82; Seaveriana, 116; Sorghi, 74, 75, 82; Thlaspeos, 41; tubulosa, 75, 82; Urbaniana, 75, 81, 82; Xanthii, 79, 82
Pucciniosira, 62; pallidula, 64, 82
Pustularia bolarioides, 4; vesiculosa, 4
Pyrenophora, 205; phaeocomes, 206; trichostoma, 206

Pyronema confluens, 4, 195
Pyrus Malus, 313; sitchensis, 238
Pythium proliferum, 168

Quamoclit, 98
Quercus, 300, 313, 315, 321; alba, 313, 318; borealis, 319; montana, 313, 318; nigra, 319; stellata, 313, 318; velutina, 319

Rand, Frederick V., Erwin F. Smith, 181
Randia mitis, 223; pubescens, 223
Ravenelia, 62; Caesalpiniae, 65, 82; Indigoferae, 65, 81; Ingae, 65, 81; Lonchocarpi, 65, 81; portoricensis, 65, 81; spinulosa, 66, 80; Whetzelii, 65
Reseda odorata, 23
Reticularia, 341; Lycoperdon, 110, 340–343, 345, 346, 348–351; sinuosa, 109
Rhabdium acutum, 163
Rhabdogloeum **abietinum,** 241; Pseudotsugae, 239
Rhabdospora **aristata,** 239; **Eucalypti,** 239
Rhipidium elongatum, 170
Rhizogone Symphoricarpi, 292
Rhizophidium, 162; ampullaceum, 159, 160; globosum, 160; pollinis, 161; sphaerocarpum, 160, 161
Rhizopus nigricans, 177
Rhytisma acerinum, 4
Ribes, 99, 175
Ricinus communis, 23
Romell, Lars, C. L. Shear, 49
Rosa, 243
Rosellinia, 86, 192, 199, 202, 211, 306, 319, 322, 326, 327, 330–336; aquila, 191, 201, 318, 326, 332, 334; Clavariae, 327, 335; Linderae, 86; prinicola, 86, 87
Rosen, H. R., Variations within a bacterial species—I. Morphologic variations, 251
Rutstroemia, 127, 129–138, 143, 146, 155; amentacea, 129; bulgarioides, 129–138; Curreyana, 129; firma, 129, 133, 137, 138; homocarpa, 129; macilenta, 129; tuberosa, 129
Rynchospora, 54; cephalotes, 54

Salix, 39, 42, 315, 320
Salvia occidentalis, 71, 82; officinalis, 23
Santo Domingo—II. Uredinales, Fungi of, Frank D. Kern, 60
Sapota Achras, 76, 82
Saprolegnia, 163; diclina, 170; dioica, 170; monilifera, 172; monoica, 159, 171; parasitica, 171
Sapromyces Reinschii, 170
Sarcoscypha, 128
Saxifraga, 42, 43
Schlegelia axillaris, 221; brachiata, 221; brachyantha, 221
Scirpus lacustris, 160
Scleria secans, 75, 82
Scleropleela, 207
Sclerotinia, 88, 89, 93, 128–135, 137–148, 151–153, 155, 191; adusta, 130; Alni, 146, 151; Ariae, 150; Aucupariae, 134; baccarum, 134, 135, 141, 145; baccata, 127; Betulae, 151; Candolleana, 127, 128, 132, 134, 135, 145; Cerasi, 134; cinerea, 142; Curreyana, 130–133; 143; Duriaeana, 131–133; filipes, 133; fructicola, 145, 148, 150, 154, 155; fructigena, 142, 148; Fuckeliana, 127, 141; Geranii, 145; heteroica, 150; Juglandis, 134; laxa, 142, 145, 148; Ledi, 150; Libertiana, 127, 129, 130, 132, 134, 145; megalospora, 134; Mespili, 134; minor, 145; Oxycocci, 134; Padi, 134; pseudotuberosa, 134; Rapula, 134; Rhododendri, 134, 140; Ricini, 145, 146; sclerotiorum, 135, 142; Smilacinae, 139, 145; Trifoliorum, 142; Tuba, 134; tuberosa, 127, 130, 132, 134, 145; Urnula, 134; Vaccinii, 141, 145
Sclerotinia, The monilioid species of, Edwin E. Honey, 127
Sclerotium bifrons, 151; Rolfsii, 22
Sclerotium Rolfsii, Notes on the parasitology of, Ruth Davis Paintin, 22
Scoleconectria tetraspora, 56
Seaver, Fred J., Doctor Arthur's rust work, 115; Studies in tropical Ascomycetes—IV. Some Hypocreales from Trinidad, 52; V. Species of Phyllachora, 214
Securidaca virgata, 224
Septogloeum **rhopaloideum,** 243
Septoria, 237, 238, 245; acicola, 244; **Boycei,** 236; **Ceanothi,** 237; Lychnidis, 237; **pinicola,** 237; **septoriopsidis,** 238; **Shepherdiae,** 238; **sitchensis,** 238; **Sonchi-arvensis,** 238; spadicea, 237
Septoriopsis, 238
Septosporium, 244
Sequoia sempervirens, 295
Sexuality and arrangement of the spores in the ascus of Neurospora sitophila, The, Marguerite S. Wilcox, 3
Shear, C. L., Lars Romell, 49; Notes

on the synonymy of some species of Hypoxylon, 83
Sida, 70, 82; spinosa, 70, 82; urens, 70, 82
Silphium, 99
Smilicina, 144; racemosa, 139
Smilax herbacea, 235; Smilacis, 235
Smith, Erwin F., Frederick V. Rand, 181
Snell, Walter H., W. G. Hutchinson and K. H. N. Newton, Temperature and moisture relations of Fomes roseus and Trametes subrosea, 276
Solenoplea, 310; microspora, 310
Solidago, 99
Sonchus, 98; arvensis, 238, 243; sonchifolia, 238
Sordaria, 211, 306, 335; fimicola, 4
Spathicarpa lanceolata, 218
Specularia, 98
Sphaeria, 189, 307, 308; concentrica, 307, 308, 310; corticata, 87; deusta, 309; discreta, 310; enteromelum, 313; globosa, 308; lutea, 310; marginata, 85; rubiformis, 85
Sphaeriales—I, Biologic studies in the, 187; II, Julian H. Miller, 305
Sphaerocarpus utricularis, 103
Sphaerostilbe, 55; longiascus, 55, 248; Musarum, 54, 55.
Sphaerulina, 198
Spirillum, 269
Spirogyra, 172; orthospira, 169, 173; Spreciana, 162
Spore formation in asci with fewer than eight spores, B. O. Dodge, 18
Sporobolus Berteronianus, 70, 82; inducus, 67, 82
Sporodinia grandis, 226
Sporophlyctis rostrata, 166
Stagonospora **Amorphae,** 235; **lapidoviridis,** 235; **Phaseoli,** 235
Stemonitis ferruginea, 111; flavogenita, 111; fusca, 111, 350; rufa, 104; splendens, 103, 111; splendens var. flaccida, 344, 346, 349, 351; uvifera, 111; varia, 113; virginiensis, 104
Stenolobium Stans, 66
Stigmaphyllon lingulatum, 71, 82
Stilbocrea intermedia, 57
Stilbum, 57
Stromatinia, 127, 131, 132, 134, 136, 138–140, 143, 145, 146, 153, 155; Paridis, 132, 145; pseudotuberosa, 131; Rapulum, 132, 138, 139, 145
Studies in tropical Ascomycetes—IV. Some Hypocreales from Trinidad, 52; V. Species of Phyllachora, Fred J. Seaver, 214
Studies on some California fungi, Lee Bonar, 292
Surinam, Myxomycetes from British Guiana and, Frank A. Gilbert, 27
Symphoricarpus, 292

Tabebuia, 63, 82
Tecoma Stans, 66, 82
Teichospora obducens, 201
Teichosporella sporadica, 201
Temperature and moisture relations of Fomes roseus and Trametes subrosea, Walter H. Snell, W. G. Hutchinson and K. H. N. Newton, 276
Tetrazygia elaeagnoides, 223
Thamnidium elegans, 177
Thamnomyces, 335, 336
Thelebolus stercoreus, 18
Thermopsis fabacea, 296; macrophylla, 295
Thuja plicata, 101, 103, 106, 108
Thurston, H. W., Jr., An interesting fern rust new to the United States, 44
Thymus vulgaris, 23
Tournefortia hirsutissima, 75, 82
Trabutia conspicua, 217; Randiae, 223; Zanthoxyli, 224
Trachyspora Alchemillae, 43
Trametes carnea, 284, 289; Feei, 281–283, 290; protracta, 277–279; subrosea, 276, 281–290
Trametes subrosea, Temperature and moisture relations of Fomes roseus and, Walter H. Snell, W. G. Hutchinson and K. H. N. Newton, 276
Tranzschelia, 62; punctata, 78, 82
Trematosphaeria, 198
Treponema pallidum, 89
Trichamphora pezizoidea, 28
Trichia, 340, 350; affinis, 112, 113; decipiens, 112; persimilis, 112, 113; scabra, 112, 113; typhoides, 103; varia, 113, 342, 346–351
Tricholaena rosea, 79, 82
Trichostigma octandrum, 74, 82
Trillium, 117–120, 122, 124, 125; erectum, 117, 118; grandiflorum, 117, 118; kamtschatikum, 117; obovatum, 117; recurvatum, 117; undulatum, 117
Trillium rust, M. F. Barrus, 117
Trinidad, Studies in tropical Ascomycetes—IV. Some Hypocreales from, Fred J. Seaver, 52
Tripsacum dactyloides, 79, 82
Triticum aestivum, 23
Triumfetta, 64, 82; semitriloba, 64, 82
Tropaeolum, 23
Tubifera stipitata, 28

Tympanopsis, 29

Ulmus, 319; americana, 242, 246
Unisexual conidia from bisexual mycelia, B. O. Dodge, 226
United States, An interesting fern rust new to the, H. W. Thurston, Jr., 44
Uredinales, Notes on arctic, J. C. Arthur, 41
Uredinia of Cronartium Comandrae and Melampsora Medusae, The, E. H. Moss, 36
Uredinopsis, 37; polypodophila, 44
Uredo, 62, 75, 76; Artocarpi, 75, 80; Bixae, 80; Bliti, 173; **bullula,** 77, 81; Chaetochloae, 69; Coccolobae, 75, 81; Cupheae, 76, 82; Dichromenae, 69; domingensis, 77; Eichorniae, 80, 82; eupatoriicola, 77; Hameliae, 76, 81; ignava, 76, 80; jatrophicola, 76, 80, 81; Panicimaximi, 67; Sapotae, 76, 82; suspecta, 77; **Toroiana,** 76, 82; Vernoniae-hookerianae, 77
Uromyces, 62, 124, 125, 168; appendiculatus, 66, 82; carnea, 42, 43; Clignyi, 74; columbianus, 66, 81; comedens, 78, 81; digitatus, 124; Dolicholi, 66, 80; gemmatus, 66, 81; Halstedii, 124; Hedysaripaniculati, 67, 81; hemisphaerica, 167; Hobsoni, 78; ignobilis, 67, 70, 82; leptodermus, 67, 81, 82; ovalis, 125; piauhyensis, 67, 82; pluriannulatus, 167; proeminens, 68, 81; Sporoboli, 70; Tricholaenae, 79, 82
Urophlyctis Alfalfae, 167; hemisphaerica, 168; Kriegeriana, 168; pluriannulatus, 167
Urtica, 295; gracilis, 295
Ustulina, 306, 309, 328, 333, 334, 335; deusta, 309; vulgaris, 308, 309

Vaccinium ,142,149, 150; ovatum, 296
Vachellia Farnesiana, 215
Valerianodes cayennense, 64, 82; jamaicense, 75, 82
Valota insularis, 75, 82
Valsa, 198, 322, 329–331
Variations within a bacterial species —I. Morphologic variations, H. R. Rosen, 251
Vaucheria sessilis, 171
Venturia inaequalis, 253
Vernonia, 98; cinerea, 76, 82; racemosa, 71, 82; Spregeliana, 69, 82
Verpa bohemica, 4, 18
Verticillium, 56
Viburnum, 315
Vicia villosa, 242
Vigna repens, 66, 82; vexillata, 223; sinensis, 23

Wedelia Ehrenbergii, 67, 82; reticulata, 67, 82
Wettsteinia, 198, 204, 205
Wilcox, Marguerite S., The sexuality and arrangement of the spores in the ascus of Neurospora sitophila, 3
Wylie, Robert B., Bruce Fink, Lichenologist, 1

Xanthium chinense, 79, 82
Xylaria, 192, 195, 199, 203, 309, 330–337; allantoidea, 332; anisopleura, 332, 334, 335; apiculata, 332; globosa, 308; obovata, 332; polymorpha, 194, 202
Xylobotryon, 333

Zanthoxylum insulare, 225; martinicense, 225
Zea everta, 23; Mays, 74, 75, 82
Zonanthemis, 99
Zygnema, 164, 173; cruciatum, 168; insigne, 163